

---

# Playing Muller Games in a Hurry

Joint work with John Fearnley, University of Warwick

Martin Zimmermann

RWTH Aachen University

June 18th, 2010

GandALF 2010  
Minori, Italy

# Motivation

---

Robert McNaughton: *Playing Infinite Games in Finite Time*. In: *A Half-Century of Automata Theory*, World Scientific (2000).

# Motivation

---

Robert McNaughton: *Playing Infinite Games in Finite Time*. In: *A Half-Century of Automata Theory*, World Scientific (2000).

*We believe that infinite games might have an interest for casual living-room recreation.*

# Motivation

---

Robert McNaughton: *Playing Infinite Games in Finite Time*. In: *A Half-Century of Automata Theory*, World Scientific (2000).

*We believe that infinite games might have an interest for casual living-room recreation.*

McNaughton suggests a method of keeping score to declare a winner such that

*.. if the play were to continue with each [player] playing forever as he has so far, then the player declared to be the winner would be the winner of the infinite play of the game.*

# Motivation

---

Robert McNaughton: *Playing Infinite Games in Finite Time*. In: *A Half-Century of Automata Theory*, World Scientific (2000).

*We believe that infinite games might have an interest for casual living-room recreation.*

McNaughton suggests a method of keeping score to declare a winner such that

*.. if the play were to continue with each [player] playing forever as he has so far, then the player declared to be the winner would be the winner of the infinite play of the game.*

“Winning regions should be equal”

# A first idea

---

Consider an infinite game  $\mathcal{G}$  played on a finite graph.

- Stop a play as soon as a cycle is closed. The winner of the induced infinite play is declared to win the finite play.
- If  $\mathcal{G}$  is positionally determined, then the winning regions of both games coincide.

# A first idea

---

Consider an infinite game  $\mathcal{G}$  played on a finite graph.

- Stop a play as soon as a cycle is closed. The winner of the induced infinite play is declared to win the finite play.
- If  $\mathcal{G}$  is positionally determined, then the winning regions of both games coincide.
- This can be extended to games  $\mathcal{G}$  that are determined with finite-state strategies: wait for a repetition of a memory state (for some fixed memory structure).

# A first idea

---

Consider an infinite game  $\mathcal{G}$  played on a finite graph.

- Stop a play as soon as a cycle is closed. The winner of the induced infinite play is declared to win the finite play.
- If  $\mathcal{G}$  is positionally determined, then the winning regions of both games coincide.
- This can be extended to games  $\mathcal{G}$  that are determined with finite-state strategies: wait for a repetition of a memory state (for some fixed memory structure).

Drawbacks (assuming  $\mathcal{G}$  is a Muller game with  $n$  vertices):

- maximal play length:  $n!$ .
- need to remember  $n!$  memory states.

**Our goal:** improve both bounds.

# Outline

---

1. Muller Games and Scoring Functions
2. Finite-time Muller Games
3. Conclusion

# Muller Games

---

- Arena:  $G = (V, V_0, V_1, E)$  with finite, directed graph  $(V, E)$ , partition  $(V_0, V_1)$  of  $V$ .

# Muller Games

---

- Arena:  $G = (V, V_0, V_1, E)$  with finite, directed graph  $(V, E)$ , partition  $(V_0, V_1)$  of  $V$ .
- Muller game:  $(G, \mathcal{F}_0, \mathcal{F}_1)$  with partition  $(\mathcal{F}_0, \mathcal{F}_1)$  of  $2^V$ .
- Player  $i$  wins play  $\rho$  iff  $\text{Inf}(\rho) = \{v \mid \exists^\omega j \text{ s.t. } \rho_j = v\} \in \mathcal{F}_i$ .

# Muller Games

---

- Arena:  $G = (V, V_0, V_1, E)$  with finite, directed graph  $(V, E)$ , partition  $(V_0, V_1)$  of  $V$ .
- Muller game:  $(G, \mathcal{F}_0, \mathcal{F}_1)$  with partition  $(\mathcal{F}_0, \mathcal{F}_1)$  of  $2^V$ .
- Player  $i$  wins play  $\rho$  iff  $\text{Inf}(\rho) = \{v \mid \exists^\omega j \text{ s.t. } \rho_j = v\} \in \mathcal{F}_i$ .
- Set of strategies for Player  $i$ :  $\Pi_i$ .
- Unique play started at  $v$  that is played according to  $\sigma \in \Pi_i$  and  $\tau \in \Pi_{1-i}$ :  $\text{Play}(v, \sigma, \tau)$ .

# Muller Games

---

- Arena:  $G = (V, V_0, V_1, E)$  with finite, directed graph  $(V, E)$ , partition  $(V_0, V_1)$  of  $V$ .
- Muller game:  $(G, \mathcal{F}_0, \mathcal{F}_1)$  with partition  $(\mathcal{F}_0, \mathcal{F}_1)$  of  $2^V$ .
- Player  $i$  wins play  $\rho$  iff  $\text{Inf}(\rho) = \{v \mid \exists^\omega j \text{ s.t. } \rho_j = v\} \in \mathcal{F}_i$ .
- Set of strategies for Player  $i$ :  $\Pi_i$ .
- Unique play started at  $v$  that is played according to  $\sigma \in \Pi_i$  and  $\tau \in \Pi_{1-i}$ :  $\text{Play}(v, \sigma, \tau)$ .
- Winning region of Player  $i$ :

$$W_i = \{v \in V \mid \exists \sigma \in \Pi_i \forall \tau \in \Pi_{1-i} : \\ \text{Play}(v, \sigma, \tau) \text{ won by Player } i\}$$

# Scoring Functions

---

For  $F \subseteq V$  define  $\text{Sc}_F: V^+ \rightarrow \mathbb{N}$ :

$$\text{Sc}_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } \text{Occ}(x_i) = F \text{ for all } i\}$$

where  $\text{Occ}(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w		a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$														

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } \text{Occ}(x_i) = F \text{ for all } i\}$$

where  $\text{Occ}(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w		a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$		0												

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w		a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$		0	0											

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1										

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1									

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w		a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$		0	0	1	1	2								

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2							

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3						

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

## Example:

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0					

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0				

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1			

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1	0		

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1	0	0	

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1	0	0	0

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

## Example:

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1	0	0	0
$Sc_{\{a,b,c\}}$	0	0	0	0	0	0	0						

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } \text{Occ}(x_i) = F \text{ for all } i\}$$

where  $\text{Occ}(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

## Example:

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1	0	0	0
$Sc_{\{a,b,c\}}$	0	0	0	0	0	0	0	1					

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } \text{Occ}(x_i) = F \text{ for all } i\}$$

where  $\text{Occ}(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

## Example:

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1	0	0	0
$Sc_{\{a,b,c\}}$	0	0	0	0	0	0	0	1	1	1			

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } \text{Occ}(x_i) = F \text{ for all } i\}$$

where  $\text{Occ}(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

## Example:

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1	0	0	0
$Sc_{\{a,b,c\}}$	0	0	0	0	0	0	0	1	1	1	2		

# Scoring Functions

---

For  $F \subseteq V$  define  $Sc_F: V^+ \rightarrow \mathbb{N}$ :

$$Sc_F(w) = \max\{k \mid \text{exist words } x_1, \dots, x_k \in V^+ \text{ s.t.} \\ x_1 \cdots x_k \text{ is suffix of } w \text{ and } Occ(x_i) = F \text{ for all } i\}$$

where  $Occ(w_1 \cdots w_n) = \{v \in V \mid \exists j \text{ s.t. } w_j = v\}$ .

## Example:

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$Sc_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1	0	0	0
$Sc_{\{a,b,c\}}$	0	0	0	0	0	0	0	1	1	1	2	2	2

# Scoring Functions

---

For  $\mathcal{F} \subseteq 2^V$  define  $\text{MaxSc}_{\mathcal{F}}: V^+ \cup V^\omega \rightarrow \mathbb{N} \cup \{\infty\}$ :

$$\text{MaxSc}_{\mathcal{F}}(\rho) = \max_{F \in \mathcal{F}} \max_{w \sqsubseteq \rho} \text{Sc}_F(w)$$

# Scoring Functions

---

For  $\mathcal{F} \subseteq 2^V$  define  $\text{MaxSc}_{\mathcal{F}}: V^+ \cup V^\omega \rightarrow \mathbb{N} \cup \{\infty\}$ :

$$\text{MaxSc}_{\mathcal{F}}(\rho) = \max_{F \in \mathcal{F}} \max_{w \sqsubseteq \rho} \text{Sc}_F(w)$$

**Example:**

w	a	a	b	b	a	a	b	c	a	b	c	a	c
$\text{Sc}_{\{a,b\}}$	0	0	1	1	2	2	3	0	0	1	0	0	0
$\text{Sc}_{\{a,b,c\}}$	0	0	0	0	0	0	0	1	1	1	2	2	2

$\mathcal{F} = \{\{a, b\}, \{a, b, c\}\}$ :

$$\text{MaxSc}_{\mathcal{F}}(w) = 3$$

# Outline

---

1. Muller Games and Scoring Functions
- 2. Finite-time Muller Games**
3. Conclusion

# Results about Scoring

---

## Lemma

Every  $w \in V^*$  with  $|w| \geq k^{|V|}$  satisfies  $\text{MaxSc}_{2^V}(w) \geq k$ .

“If you play long enough, some score value will be high”

Lower bound: there are words  $w_k$  of length  $k^{|V|} - 1$  with  $\text{MaxSc}_{2^V}(w_k) < k$ .

# Results about Scoring

---

## Lemma

Every  $w \in V^*$  with  $|w| \geq k^{|V|}$  satisfies  $\text{MaxSc}_{2V}(w) \geq k$ .

“If you play long enough, some score value will be high”

Lower bound: there are words  $w_k$  of length  $k^{|V|} - 1$  with  $\text{MaxSc}_{2V}(w_k) < k$ .

## Lemma (McNaughton 2000)

Let  $k, \ell \geq 2$ , let  $F, H \subseteq V$ , let  $w \in V^*$  and  $s \in V$  such that  $\text{Sc}_F(w) < k$  and  $\text{Sc}_H(w) < \ell$ . If  $\text{Sc}_F(ws) = k$  and  $\text{Sc}_H(ws) = \ell$ , then  $F = H$ .

“At most one score value can increase at a time”

# Finite-time Muller Games

---

- Finite-time Muller game:  $(G, \mathcal{F}_0, \mathcal{F}_1, k)$  with threshold  $k \geq 2$ .
- Play: path  $w = w_1 \cdots w_n$  with  $\text{MaxSc}_{2^V}(w_0 \cdots w_n) = k$ , but  $\text{MaxSc}_{2^V}(w_1 \cdots w_{n-1}) < k$ .
- Previous Lemma yields unique  $F \subseteq V$  such that  $\text{Sc}_F(w) = k$ . Player  $i$  wins  $w$  iff  $F \in \mathcal{F}_i$ .
- Strategies and winning regions defined as usual.

# Finite-time Muller Games

---

- Finite-time Muller game:  $(G, \mathcal{F}_0, \mathcal{F}_1, k)$  with threshold  $k \geq 2$ .
- Play: path  $w = w_1 \cdots w_n$  with  $\text{MaxSc}_{2^V}(w_0 \cdots w_n) = k$ , but  $\text{MaxSc}_{2^V}(w_1 \cdots w_{n-1}) < k$ .
- Previous Lemma yields unique  $F \subseteq V$  such that  $\text{Sc}_F(w) = k$ . Player  $i$  wins  $w$  iff  $F \in \mathcal{F}_i$ .
- Strategies and winning regions defined as usual.

McNaughton considered a different definition of a finite-time Muller game: stop play when some  $\text{Sc}_F$  reaches  $|F|! + 1$ .

## Theorem (McNaughton 2000)

*The winning regions in a Muller game and in McNaughton's finite-time Muller game coincide.*

# Main Theorem

---

## Theorem

*The winning regions in a Muller game  $(G, \mathcal{F}_0, \mathcal{F}_1)$  and in the finite-time Muller game  $(G, \mathcal{F}_0, \mathcal{F}_1, 3)$  coincide.*

# Main Theorem

---

## Theorem

*The winning regions in a Muller game  $(G, \mathcal{F}_0, \mathcal{F}_1)$  and in the finite-time Muller game  $(G, \mathcal{F}_0, \mathcal{F}_1, 3)$  coincide.*

We prove a stronger statement about winning strategies in the infinite-duration Muller game, which implies the theorem.

## Lemma

*Player  $i$  has a strategy  $\sigma$  for a Muller game  $(G, \mathcal{F}_0, \mathcal{F}_1)$  such that  $\text{MaxSc}_{\mathcal{F}_{1-i}}(\text{Play}(v, \sigma, \tau)) \leq 2$  for every  $v \in W_i$  and every  $\tau \in \Pi_{1-i}$ .*

## What about 2?

---

The bound 2 in the lemma is optimal:

Player 0 has a winning strategy, but cannot avoid score values of 2 for Player 1.



- $\mathcal{F}_0 = \{\{1, 2, 3\}, \{1\}, \{3\}\}$
- $\mathcal{F}_1 = 2^{\{1,2,3\}} \setminus \mathcal{F}_0$

One of the plays 2112 or 2332 is consistent with every winning strategy.

## What about 2?

---

The bound 2 in the lemma is optimal:

Player 0 has a winning strategy, but cannot avoid score values of 2 for Player 1.



- $\mathcal{F}_0 = \{\{1, 2, 3\}, \{1\}, \{3\}\}$
- $\mathcal{F}_1 = 2^{\{1,2,3\}} \setminus \mathcal{F}_0$

One of the plays 2112 or 2332 is consistent with every winning strategy.

Consequence:

To show that the finite-time Muller game with threshold 2 is equivalent, we need other proof techniques.

# Outline

---

1. Muller Games and Scoring Functions
2. Finite-time Muller Games
- 3. Conclusion**

# Conclusion

---

We have presented a finite-duration version of a Muller game that is equivalent to the original game.

- Reachability game on a tree; hence, simple algorithms are available.
- Our strategies are eager: they do not spend more time in “bad” loops than they have to.

# Conclusion

---

We have presented a finite-duration version of a Muller game that is equivalent to the original game.

- Reachability game on a tree; hence, simple algorithms are available.
- Our strategies are eager: they do not spend more time in “bad” loops than they have to.

	Reduction	McNaughton	here
Threshold	–	$ F ! + 1$	3
Play Length	$\leq n \cdot n! + 1$	$\leq (n! + 1)^n$	$\leq 3^n$
Space	$\mathcal{O}(n!)$	$\mathcal{O}((n! + 1)^n)$	$\mathcal{O}(3^n)$

# Open Questions

---

- Is the finite-time Muller game with threshold 2 equivalent to the original Muller game?
- Given a winning strategy for a finite-time Muller game, can we turn it into a winning strategy for the Muller game?